

**PRODUCT INFORMATION**

<b>Common Name</b>	KHK4083, KHK-4083, KHK 4083
<b>Conjugate</b>	Unconjugated
<b>Synonyms</b>	TNFRSF4;OX40;CD134;OX40L receptor;ACT35;TXGP1L
<b>Applications</b>	Flow Cyt
<b>Endotoxin</b>	Less than 1.0 EU/μg by the LAL method. For <1 EU/mg requirements, please contact us for customization.
<b>Recommended Dilutions</b>	Flow Cyt 1:100
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Host Species</b>	Homo sapiens
<b>IgG type</b>	Human IgG1(E356D,M358L) - kappa
<b>Reactivity</b>	Human
<b>Target</b>	OX40
<b>Uniprot ID</b>	P43489
<b>Description</b>	Anti-OX40(rocatinlimab biosimilar) mAb
<b>Delivery</b>	In Stock
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.
<b>Background</b>	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
<b>Usage</b>	Research use only






Figure 1. Flow cytometry analysis with 0.5 µg/ml Anti-OX40(rocatinlimab biosimilar) mAb (BME100654) on HEK293 cells transfected with human OX40 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

