

**PRODUCT INFORMATION**

<b>Common Name</b>	MEDI0562
<b>Conjugate</b>	Unconjugated
<b>Synonyms</b>	TNFRSF4, TXGP1L, CD134, ACT35 antigen, OX40L receptor
<b>Applications</b>	ELISA, Flow Cyt
<b>Recommended Dilutions</b>	ELISA 1:5000-10000, Flow Cyt 1:100
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Host Species</b>	Chimeric Humanized
<b>IgG type</b>	Human IgG1 - kappa
<b>Reactivity</b>	Human
<b>Target</b>	OX40
<b>Uniprot ID</b>	P43489
<b>Description</b>	Anti-OX40(tavolimab biosimilar) mAb
<b>Delivery</b>	In Stock
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.
<b>Background</b>	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
<b>Usage</b>	Research use only






Figure 1. Flow cytometry analysis with 0.5 µg/mL Anti-OX40(tavolimab biosimilar) mAb (BME100047) on HEK293 cells transfected with Human OX40 protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

### Anti-OX40(tavolimab biosimilar)mAb ELISA

0.2 µg of Human OX40, hFc-His tagged protein per well




Figure 2. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human OX40 Protein, hFc-His Tag (PME100022) can bind Anti-OX40(tavolimab biosimilar)mAb (BME100047) in a linear range of 0.76-20.58 ng/mL.

