

PRODUCT INFORMATION

Common Name	GBR-830
Conjugate	Unconjugated
Synonyms	TNFRSF4;OX40;CD134;OX40L receptor;ACT35;TXGP1L
Applications	ELISA, Flow Cyt
Endotoxin	Less than 1.0 EU/ μ g by the LAL method. For <1 EU/mg requirements, please contact us for customization.
Recommended Dilutions	ELISA 1:5000-10000, Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Humanized
IgG type	Human IgG1(E356D,M358L) - kappa
Reactivity	Human
Target	OX40
Uniprot ID	P43489
Description	Anti-OX40(telazorlimab biosimilar) mAb
Delivery	In Stock
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 μ m) prior to use.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only



Anti-OX40(telazorlimab biosimilar) mAb ELISA

0.2 μ g of Human OX40, hFc-His tagged protein per well




Figure 1. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human OX40 Protein, hFc-His tag(PME100022) can bind Anti-OX40(telazorlimab biosimilar) mAb (BME100692) in a linear range of 8-40 ng/mL. In order to specifically detect BME100692, mouse anti-human Fab-specific antibody was used as detection antibody.




Figure 2. Flow cytometry analysis with 0.5 μ g/ml Anti-OX40(telazorlimab biosimilar) mAb (BME100692) on HEK293 cells transfected with Human OX40 protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).



Anti-OX40 (telazorlimab biosimilar) mAb ELISA

0.2 μ g of Cynomolgus OX40, hFc tagged protein per well




Figure 3. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Cynomolgus OX40 Protein, hFc Tag (PME-C100100) can bind Anti-OX40(telazorlimab biosimilar) mAb (BME100692) in a linear range of 8-40 ng/mL.

