


**PRODUCT INFORMATION**

<b>Clone ID</b>	DM42
<b>Target</b>	S protein
<b>Synonyms</b>	SARS-CoV-2 Spike S2
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-SARS-CoV-2 Spike antibody(DM42); Rabbit mAb
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P0DTC2
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	SARS-CoV-2
<b>Applications</b>	ELISA
<b>Recommended Dilutions</b>	ELISA 1:5000-10000
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Endotoxin</b>	Less than 1.0 EU/μg by the LAL method. For <1 EU/mg requirements, please contact us for customization.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) also known as Covid19 (2019 Novel Coronavirus) is a virus that causes illnesses ranging from the common cold to severe diseases. The spike protein is a type I transmembrane protein containing two subunits; S1 and S2. S1 mainly contains a receptor binding domain (RBD); which accounts for recognizing the cell surface receptor; ACE2. S2 contains basic elements needed for the membrane fusion. Recent publications indicate that S1-RBD domain can induce virus neutralizing-antibody and T cell response.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated
<b>DIMA Disclaimer</b>	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr





**Figure 1.** Elisa plate pre-coated by 2 µg/ml(100µl/well) SARS-CoV-2 Spike S2 protein can bind Rabbit Anti-SARS-CoV-2 Spike S2 monoclonal antibody (**clone:DM42**) in a linear range of 0.32-40 ng/ml.

DIMABIO CONFIDENTIAL

