

PRODUCT INFORMATION

Clone ID	3E11
Target	SIGLEC10
Synonyms	SIGLEC10;Siglec-10;SLG2;PRO940;MGC126774
Host Species	Rabbit
Description	Anti-SIGLEC10 antibody(3E11), IgG1 Chimeric mAb
Delivery	In Stock
Uniprot ID	Q96LC7
IgG type	Rabbit/Human Fc chimeric IgG1
Clonality	Monoclonal
Reactivity	Human
Applications	ELISA, Flow Cyt
Recommended Dilutions	ELISA 1/5000-10000;Flow Cyt 1/100
Purification	Purified from cell culture supernatant by affinity chromatography
Endotoxin	Less than 1.0 EU/μg by the LAL method. For <1 EU/mg requirements, please contact us for customization.
Formulation & Reconstitution	Powder
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 μm) prior to use.
Background	SIGLECs are members of the immunoglobulin superfamily that are expressed on the cell surface. Most SIGLECs have 1 or more cytoplasmic immune receptor tyrosine-based inhibitory motifs, or ITIMs. SIGLECs are typically expressed on cells of the innate immune system, with the exception of the B-cell expressed SIGLEC6 (MIM 604405).
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Flow cytometry analysis with 10 μ g/mL Anti-SIGLEC10 antibody(3E11) mAb on HEK293 cells transfected with human SIGLEC10 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Anti-SIGLEC10 antibody(3E11), IgG1 Chimeric mAb ELISA
 0.1 μ g of Human SIGLEC10, His tagged protein per well




Figure 2. ELISA plate pre-coated by 1 μ g/ml (100 μ l/well) Human SIGLEC10 Protein, His Tag can bind IgG1 Chimeric anti-SIGLEC10 monoclonal antibody(clone: 3E11) in a linear range of 40-10000 ng/ml.

