

PRODUCT INFORMATION

Common Name	PR-1808164, ABBV-0805, BAN-0805, PD-0805
Conjugate	Unconjugated
Synonyms	SYUA, NACP, PARK1
Applications	Flow Cyt
Endotoxin	Less than 1.0 EU/μg by the LAL method. For <1 EU/mg requirements, please contact us for customization.
Recommended Dilutions	Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Humanized, Chimeric
IgG type	Human IgG4 - Kappa
Reactivity	Human
Target	SNCA
Uniprot ID	P37840
Description	Anti-SNCA(exidavnemab biosimilar) mAb
Delivery	In Stock
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 μm) prior to use.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only






Figure 1. Flow cytometry analysis with 0.5 $\mu\text{g}/\text{mL}$ Anti-SNCA(exidavnemab biosimilar) mAb (BME100503) on HEK293 cells transfected with SNCA protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

