

PRODUCT INFORMATION

Common Name	JR-141
Conjugate	Unconjugated
Synonyms	TR;TfR;TfR1;Trfr;T9;p90;CD71
Applications	ELISA, Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000, Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Humanized
IgG type	Human IgG1 - kappa
Reactivity	Human
Target	TFRC
Uniprot ID	P02786
Description	Anti-TFRC(pabinafusp alfa biosimilar) mAb
Delivery	In Stock
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.






Figure 1. Flow cytometry analysis with 1  $\mu$ g/mL Anti-TFRC mAb (BME100138) on Expi293 cells transfected with Human TFRC protein (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).

**Anti-TFRC(pabinafusp alfa biosimilar) mAb ELISA**  
0.2  $\mu$ g of Human TFRC, His tagged protein per well




Figure 2. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human TFRC Protein, His Tag (PME100775) can Anti-TFRC(pabinafusp alfa biosimilar) mAb (BME100138) in a linear range of 0.13-16 ng/mL.






Figure 3. Flow cytometry analysis of antigen binding of anti-human TFRC mAb(BME100138).  
(A) BME100138 does not bind to CHO-S cells that do not express TFRC.  
(B) A clear peak shift of BME100138 was seen compared to the control when incubated with TFRC-expressing Siha cells, indicating strong binding of BME100138 to TFRC. Antibodies were incubated at 5 µg/mL.

