

PRODUCT INFORMATION

Clone ID	DMC467
Target	TGFBR2
Synonyms	AAT3; FAA3; LDS1B; LDS2; LDS2B; MFS2; RIIC; TAAD2; TBR-ii; TBRll; TGFbeta-Rll; TGFR-2
Host Species	Rabbit
Description	Anti-TGFBR2 antibody(DMC467); IgG1 Chimeric mAb
Delivery	In Stock
Uniprot ID	P37173
IgG type	Rabbit/Human Fc chimeric IgG1
Clonality	Monoclonal
Reactivity	Human
Applications	Flow Cyt
Recommended Dilutions	Flow Cyt 1:100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene is a transmembrane protein that has a protein kinase domain; forms a heterodimeric complex with TGF-beta receptor type-1; and binds TGF-beta. This receptor:ligand complex phosphorylates proteins; which then enter the nucleus and regulate the transcription of genes related to cell proliferation; cell cycle arrest; wound healing; immunosuppression; and tumorigenesis. Mutations in this gene have been associated with Marfan Syndrome; Loeys-Deitz Aortic Aneurysm Syndrome; and the development of various types of tumors. Alternatively spliced transcript variants encoding different isoforms have been characterized. [provided by RefSeq; Aug 2017]
Usage	Research use only
Conjugate	Unconjugated
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr






Figure 1. Flow cytometry analysis with Anti-TGFBR2 (DMC467) on HEK293 cells transfected with human TGFR2 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).




Figure 2. Flow cytometry analysis of antigen binding of anti-human TGFR2 mAb(DMC100467).

(A) DMC100467 does not bind to CHO-S cells that do not express TGFR2.

(B) A clear peak shift of DMC100467 was seen compared to the control when incubated with TGFR2-expressing hepG2 cells, indicating strong binding of DMC100467 to TGFR2. Antibodies were incubated at 5 μ g/ml.

