


**PRODUCT INFORMATION**

<b>Common Name</b>	MTIG-7192-A, MTIG-7192A, MTIG7192A, RG-6058, RO-7092284, RO7092284
<b>Conjugate</b>	Unconjugated
<b>Synonyms</b>	TIGIT;SIG9;STM3
<b>Applications</b>	ELISA; Flow Cyt
<b>Recommended Dilutions</b>	ELISA 1:5000-10000; Flow Cyt 1:100
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Host Species</b>	Homo sapiens
<b>IgG type</b>	Human IgG1 - kappa
<b>Reactivity</b>	Human
<b>Target</b>	TIGIT
<b>Uniprot ID</b>	Q495A1
<b>Description</b>	Anti-TIGIT (tiragolumab biosimilar) mAb
<b>Delivery</b>	In Stock
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
<b>Usage</b>	Research use only




## Anti-TIGIT (tiragolumab biosimilar) mAb ELISA

0.2  $\mu$ g of Human TIGIT, His tagged protein per well



**Figure 1.** ELISA plate pre-coated by 2  $\mu$ g/ml (100  $\mu$ l/well) Human TIGIT, His tagged protein ([getskuurl sku="PME100506"]) can bind Anti-TIGIT Neutralizing antibody in a linear range of 1.6-200.0 ng/ml.



**Figure 2.** Flow cytometry analysis with 15  $\mu$ g/mL Anti-TIGIT (tiragolumab) mAb (BME100026) on HEK293 cells transfected with Human TIGIT protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

