



PRODUCT INFORMATION

Clone ID	DM141
Target	TNFSF12
Synonyms	TNFSF12
Host Species	Rabbit
Description	Anti-TNFSF12 antibody(DM141); Rabbit mAb
Delivery	In Stock
Uniprot ID	O43508
IgG type	Rabbit IgG
Clonality	Monoclonal
Reactivity	Human
Applications	ELISA; Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000; Flow Cyt 1:100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This protein is a ligand for the FN14:TWEAKR receptor. This cytokine has overlapping signaling functions with TNF; but displays a much wider tissue distribution. This cytokine; which exists in both membrane-bound and secreted forms; can induce apoptosis via multiple pathways of cell death in a cell type-specific manner. This cytokine is also found to promote proliferation and migration of endothelial cells; and thus acts as a regulator of angiogenesis. Alternative splicing results in multiple transcript variants. Some transcripts skip the last exon of this gene and continue into the second exon of the neighboring TNFSF13 gene; such read-through transcripts are contained in GeneID 407977; TNFSF12-TNFSF13.
Usage	Research use only
Conjugate	Unconjugated
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.





**Figure 1.** ELISA plate pre-coated by 1 µg/ml (100 µl/well) Human TNFSF12 protein, hFc tagged protein ([getskuurl sku="PME100105"]) can bind Rabbit anti-TNFSF12 monoclonal antibody(**clone: DM141**) in a linear range of 0. 1-10 ng/ml.



**Figure 2.** Flow cytometry analysis with Anti-TNFSF12 (DM141) on HEK293 cells transfected with human TNFSF12 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

