

## **PRODUCT INFORMATION**

**Clone ID** 17A11 **Target** Trop2

TACSTD2;GA733-1;M1S1;TROP2 **Synonyms** 

**Host Species** Rabbit

Description Anti-TROP2 antibody(17A11), Rabbit mAb

**Delivery** In Stock P09758 **Uniprot ID** IgG type Rabbit IgG Clonality Monoclonal Reactivity Human **Applications** WB

Recommended WB 1:1000 **Dilutions** 

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

This intronless gene encodes a carcinoma-associated antigen. This antigen is a cell surface

receptor that transduces calcium signals.

Mutations of this gene have been associated with gelatinous drop-like corneal dystrophy.

Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only

**Background** 





130 kDa-

100 kDa-

70 kDa-

55 kDa-

35 kDa-

25 kDa-

Figure 1.Anti-Trop2 antibody (SKU# DME101001) at 1/1000 dilution

Lane: A431, whole cell lysate

Secondary: Goat Anti-Rabbit IgG H&L (HRP) at 1/5000 dilution

Predicted band size: 36 kDa Observed band size: 40 kDa



Email: info@dimabio.com Website: www.dimabio.com