Cat. No. DMC100465B



## **PRODUCT INFORMATION**

Clone ID **DMC465** CCR1 **Target** 

CD191; CKR-1; CKR1; CMKBR1; HM145; MIP1aR; **Synonyms** 

SCYAR1

**Host Species** 

Biotinylated Anti-CCR1 antibody(DMC465); IgG1 **Description** 

Chimeric mAb

Delivery 2-3 weeks P32246 **Uniprot ID** 

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human

**Applications** Flow Cyt; WB

Recommended **Dilutions** 

Flow Cyt 1:100; WB 1:1000

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Formulation & Reconstitution

**Background** 

Storage & Shipping

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a member of the beta chemokine receptor family; which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. The ligands of this receptor include macrophage inflammatory protein 1 alpha (MIP-1 alpha); regulated on activation normal T expressed and secreted protein (RANTES); monocyte chemoattractant protein 3 (MCP-3); and myeloid progenitor inhibitory factor-1 (MPIF-1). Chemokines and their

receptors mediated signal transduction are critical for the recruitment of effector immune cells to the site of inflammation. Knockout studies of the mouse homolog suggested the roles of this gene in host protection from inflammatory response; and susceptibility to virus and parasite. This gene and other chemokine receptor genes; including CCR2; CCRL2; CCR3; CCR5 and CCXCR1;

are found to form a gene cluster on chromosome 3p. [provided by RefSeq; Jul 2008]

Usage Research use only

Conjugate Biotinylated

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are

> > Email: info@dimabio.com

Website: www.dimabio.com

actively scrutinizing all patent application to

ensure no IP infringement.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

**DIMA Disclaimer** 

