

## **PRODUCT INFORMATION**

Clone ID	DMC270	
Target	CD160	
Synonyms	BY55; NK1; NK28	
Host Species	Rabbit	
Description	Biotinylated Anti-CD160 antibody(DMC270); IgG1 Chimeric mAb	
Delivery	2-3 weeks	
Uniprot ID	095971	
lgG type	Rabbit/Human Fc chimeric IgG1	
Clonality	Monoclonal	
Reactivity	Human	
Applications	Flow Cyt	
Recommended Dilutions	Flow Cyt 1:100	
Purification	Purified from cell culture supernatant by affinity chromatography	
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.	
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.	
Background	CD160 is an 27 kDa glycoprotein which was initially identified with the monoclonal antibody BY55. Its expression is tightly associated with peripheral blood NK cells and CD8 T lymphocytes with cytolytic effector activity. The cDNA sequence of CD160 predicts a cysteine-rich; glycosylphosphatidylinositol-anchored protein of 181 amino acids with a single Ig-like domain weakly homologous to KIR2DL4 molecule. CD160 is expressed at the cell surface as a tightly disulfide-linked multimer. RNA blot analysis revealed CD160 mRNAs of 1.5 and 1.6 kb whose expression was highly restricted to circulating NK and T cells; spleen and small intestine. Within NK cells CD160 is expressed by CD56dimCD16 cells whereas among circulating T cells its expression is mainly restricted to TCRgd bearing cells and to TCRab CD8brightCD95 CD56 CD28-CD27-cells. In tissues; CD160 is expressed on all intestinal intraepithelial lymphocytes. CD160 shows a broad specificity for binding to both classical and nonclassical MHC class I molecules.	
Usage	Research use only	
Conjugate	Biotinylated	
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.	
ddress: Wuhan institute of iotechnology B7, Biolake N aoxin Road, Wuhan, Hube alaphone: +1 2400940618	f Email: info@dimabio.com No.666 Website: www.dimabio.com i, China	

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)



Cat. No. DMC100270B



MARIO CONTRACTION

Email: info@dimabio.com Website: www.dimabio.com

