

**PRODUCT INFORMATION**

|   |  |
|---|--|
| <b>Clone ID</b>                         | DM30   |
| <b>Target</b>                           | CD38   |
| <b>Synonyms</b>                         | T10; cADPr hydrolase 1   |
| <b>Host Species</b>                     | Rabbit   |
| <b>Description</b>                      | Biotinylated Anti-CD38 antibody(DM30); Rabbit mAb  |
| <b>Delivery</b>                         | 2-3 weeks  |
| <b>Uniprot ID</b>                       | P28907   |
| <b>IgG type</b>                         | Rabbit IgG   |
| <b>Clonality</b>                        | Monoclonal   |
| <b>Reactivity</b>                       | Human  |
| <b>Applications</b>                     | ELISA; Flow Cyt  |
| <b>Recommended Dilutions</b>            | ELISA 1:5000-10000; Flow Cyt 1:100   |
| <b>Purification</b>                     | Purified from cell culture supernatant by affinity chromatography  |
| <b>Endotoxin</b>                        | Less than 1.0 EU/μg by the LAL method. For <1 EU/mg requirements, please contact us for customization.   |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.   |
| <b>Storage&amp;Shipping</b>             | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.  |
| <b>Sterility</b>                        | Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 μm) prior to use.  |
| <b>Background</b>                       | CD antigen CD38 is also known as ADP-ribosyl cyclase 1; which belongs to the ADP-ribosyl cyclase family. CD38 is expressed at high levels in pancreas; liver; kidney; brain; testis; ovary; placenta; malignant lymphoma and neuroblastoma. CD38 is a multifunctional ectoenzyme that catalyzes the synthesis and hydrolysis of cyclic ADP-ribose (cADPR) from NAD to ADP-ribose. These reaction products are essential for the regulation of intracellular Ca <sup>2+</sup> . The loss of CD38 function is associated with impaired immune responses; metabolic disturbances; and behavioral modifications. The CD38 protein is a marker of cell activation. It has been connected to HIV infection; leukemias; myelomas; solid tumors; type II diabetes mellitus and bone metabolism. CD38 has been used as a prognostic marker in leukemia. |
| <b>Usage</b>                            | Research use only  |
| <b>Conjugate</b>                        | Biotinylated   |
| <b>DIMA Disclaimer</b>                  | All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr   |



DIMABIO CONFIDENTIAL

