


PRODUCT INFORMATION

Common Name	INX 021;INX021;INX-021;SAR-441344;SAR441344;SAR 441344
Conjugate	Biotinylated
Synonyms	IGM;IMD3;TRAP;gp39;CD154;CD40L;HIGM1;T-BAM;TNFSF5;hCD40L
Applications	ELISA
Endotoxin	Less than 1.0 EU/μg by the LAL method. For <1 EU/mg requirements, please contact us for customization.
Recommended Dilutions	ELISA 1:5000-10000
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	N/A
IgG type	IgG1
Reactivity	Human
Target	CD40 Ligand
Uniprot ID	P29965
Description	Biotinylated Anti-CD40 Ligand(frexalimab biosimilar) mAb
Delivery	In Stock
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 μm) prior to use.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only



Biotinylated Anti-CD40 Ligand(frexalimab biosimilar) ELISA

0.2 μ g of Human CD40 Ligand, hFc tagged protein per well



Biotinylated Anti-CD40 Ligand(frexalimab biosimilar) mAb (ng/mL)

Figure 1. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human CD40 Ligand Protein, hFc tag (PME100192) can bind Biotinylated Anti-CD40 Ligand(frexalimab biosimilar) mAb (BME100516B) in a linear range of 80-400 ng/mL .

