

## PRODUCT INFORMATION

<b>Clone ID</b>	DMC687
<b>Target</b>	CXADR
<b>Synonyms</b>	CAR; CAR4:6; HCAR
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-CXADR antibody(DMC687); IgG1 Chimeric mAb
<b>Delivery</b>	2-3 weeks
<b>Uniprot ID</b>	P78310
<b>IgG type</b>	Rabbit/Human Fc chimeric IgG1
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	Flow Cyt; WB; IHC
<b>Recommended Dilutions</b>	Flow Cyt 1:100; WB 1:1000; IHC 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein encoded by this gene is a type I membrane receptor for group B coxsackieviruses and subgroup C adenoviruses. Several transcript variants encoding different isoforms have been found for this gene. Pseudogenes of this gene are found on chromosomes 15; 18; and 21. [provided by RefSeq; May 2011]
<b>Usage</b>	Research use only

