Cat. No. DME100208B



PRODUCT INFORMATION

Clone ID DM208 **Target** CXCR3

CD182; CD183; CKR-L2; CMKAR3; GPR9; IP10-R; **Synonyms**

Mig-R; MigR

Host Species

Biotinylated Anti-CXCR3 antibody(DM208); Rabbit **Description**

mAb

Delivery 2-3 weeks **Uniprot ID** P49682 IgG type Rabbit IgG Clonality Monoclonal Reactivity Human

Applications ELISA; Flow Cyt

Recommended

Background

ELISA 1:5000-10000; Flow Cyt 1:100 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation &

Reconstitution for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a G protein-coupled receptor with selectivity for three chemokines; termed CXCL9:Mig (monokine induced by interferon-g); CXCL10:IP10 (interferon-g-inducible 10 kDa protein) and CXCL11:I-TAC (interferon-inducible T cell a-chemoattractant). Binding of chemokines to this protein induces cellular responses that are involved in leukocyte traffic; most notably

integrin activation; cytoskeletal changes and chemotactic migration. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. One of the isoforms (CXCR3-B) shows high affinity binding to chemokine; CXCL4:PF4 (PMID:12782716).

[provided by RefSeq; Jun 2011]

Usage Research use only

Conjugate Biotinylated

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are

> > Email: info@dimabio.com Website: www.dimabio.com

actively scrutinizing all patent application to

ensure no IP infringement.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)

DIMA Disclaimer

