

## PRODUCT INFORMATION

<b>Clone ID</b>	DMC478
<b>Target</b>	GPR87
<b>Synonyms</b>	FKSG78; GPR95; KPG_002
<b>Host Species</b>	Rabbit
<b>Description</b>	Biotinylated Anti-GPR87 antibody(DMC478); IgG1 Chimeric mAb
<b>Uniprot ID</b>	Q9BY21
<b>IgG type</b>	Rabbit/Human Fc chimeric IgG1
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	Flow Cyt
<b>Recommended Dilutions</b>	Flow Cyt 15µg/ml
<b>Delivery</b>	In Stock
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-coupled receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in lung squamous cell carcinoma (PMID:18057535) and regulated by p53 (PMID:19602589). [provided by RefSeq; Nov 2011]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Biotinylated
<b>DIMA Disclaimer</b>	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr






Figure 1. Flow cytometry analysis with 15µg/mL Biotinylated Anti-GPR87 (DMC478) on HEK293 cells transfected with human GPR87 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

