

PRODUCT INFORMATION

Clone ID DMC271 **Target** II 1B

Synonyms IL-1; IL1-BETA; IL1F2

Host Species Rabbit

Description Anti-IL1B antibody(DMC271); IgG1 Chimeric mAb

Delivery 2-3 weeks **Uniprot ID** P01584

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended

Dilutions

Flow Cyt 1:100

Purified from cell culture supernatant by affinity **Purification**

chromatography

Formulation & Reconstitution

Background

Storage & Shipping

Lyophilized from sterile PBS, pH 7.4. Normally 5 % 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine is produced by activated macrophages as a proprotein; which is proteolytically processed to its active form by caspase 1 (CASP1:ICE). This cytokine is an important mediator of the inflammatory response; and is involved in a

variety of cellular activities; including cell proliferation; differentiation; and apoptosis. The induction of cyclooxygenase-2 (PTGS2:COX2) by this cytokine in the central nervous system (CNS) is found to contribute to inflammatory pain hypersensitivity. Similarly; IL-1B has been implicated in human osteoarthritis pathogenesis. Patients with severe Coronavirus Disease 2019

(COVID-19) present elevated levels of pro-

inflammatory cytokines such as IL-1B in bronchial alveolar lavage fluid samples. The lung damage induced by the Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is to a large extent; a result of the inflammatory response promoted by cytokines such as IL-1B. This gene and eight other interleukin 1 family

> Email: info@dimabio.com Website: www.dimabio.com

genes form a cytokine gene cluster on

chromosome 2

Usage Research use only

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

