


PRODUCT INFORMATION

Clone ID	8C4
Target	MMAE
Synonyms	N.A.
Host Species	Rabbit
Description	Biotinylated Anti-MMAE antibody(8C4); Rabbit mAb
Delivery	In Stock
Uniprot ID	N.A.
IgG type	Rabbit IgG
Clonality	Monoclonal
Reactivity	N.A.
Applications	ELISA
Recommended Dilutions	ELISA 1:5000-10000
Purification	Purified from cell culture supernatant by affinity chromatography
Endotoxin	Less than 1.0 EU/μg by the LAL method. For <1 EU/mg requirements, please contact us for customization.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	N.A.
Usage	Research use only
Conjugate	Biotinylated
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr



Biotinylated Anti-MMAE antibody(8C4); Rabbit mAb ELISA

0.2 μg of IgG-MMAE per well



Biotinylated Anti-MMAE antibody(8C4); Rabbit mAb (ng/mL)

Figure 1. ELISA plate pre-coated by 2 $\mu\text{g}/\text{mL}$ (100 $\mu\text{L}/\text{well}$) IgG-MMAE can bind Biotinylated Anti-MMAE antibody(8C4); Rabbit mAb (DME101004B) in a linear range of 370-1111 ng/mL. In order to specifically detect DME101004B, HRP Conjugated-Streptavidin was used as detection antibody.

