Cat. No. DME100123B



## **PRODUCT INFORMATION**

Clone ID DM123 PDL1 **Target** 

PD-L1; CD274; B7-H1; PDCD1L1; PDCD1LG1 **Synonyms** 

**Host Species** Rabbit

Description Anti-PD-L1 antibody(DM123); Rabbit mAb

**Delivery** 2-3 weeks **Uniprot ID** Q9NZQ7 IgG type Rabbit IgG Clonality Monoclonal Reactivity Human

**Applications** ELISA; Flow Cyt

Recommended

Storage & Shipping

**Background** 

ELISA 1:5000-10000; Flow Cyt 1:100 **Dilutions** 

Purified from cell culture supernatant by affinity Purification

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & 8% trehalose is added as protectants before Reconstitution

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

This gene encodes an immune inhibitory receptor ligand that is expressed by hematopoietic and non-hematopoietic cells; such as T cells and B cells and various types of tumor cells. The encoded protein is a type I transmembrane protein that has immunoglobulin V-like and C-like

domains. Interaction of this ligand with its receptor inhibits T-cell activation and cytokine production. During infection or inflammation of normal tissue; this interaction is important for preventing autoimmunity by maintaining homeostasis of the immune response. In tumor

microenvironments; this interaction provides an immune escape for tumor cells through cytotoxic T-cell inactivation. Expression of this gene in tumor cells is considered to be prognostic in many types of human malignancies; including colon cancer and renal cell carcinoma. Alternative splicing results in multiple transcript variants.

> Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only

