

## **PRODUCT INFORMATION**

Clone ID DM140 TNFSF12 **Target** TNFSF12 **Synonyms Host Species** Rabbit

Description Anti-TNFSF12 antibody(DM140); Rabbit mAb

**Delivery** 2-3 weeks **Uniprot ID** 043508 IgG type Rabbit IgG Clonality Monoclonal Reactivity Human

**Applications** ELISA; Flow Cyt

Recommended **Dilutions** 

ELISA 1:5000-10000; Flow Cyt 1:100

Purified from cell culture supernatant by affinity Purification

chromatography

Formulation & Reconstitution

**Background** 

Storage & Shipping

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This protein is a ligand for the FN14:TWEAKR receptor. This cytokine has overlapping signaling functions with TNF; but displays a much wider tissue distribution. This cytokine; which exists in both membrane-bound and secreted forms; can induce apoptosis via multiple pathways of cell death in a cell typespecific manner. This cytokine is also found to

promote proliferation and migration of endothelial

cells; and thus acts as a regulator of angiogenesis. Alternative splicing results in multiple transcript variants. Some transcripts skip the last exon of this gene and continue into the second exon of the neighboring TNFSF13 gene; such read-through transcripts are contained in GeneID 407977; TNFSF12-TNFSF13.

> Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only

