

## PRODUCT INFORMATION

|   |   |
|---|---|
| <b>Clone ID</b>                         | DMC464  |
| <b>Target</b>                           | TNFSF15   |
| <b>Synonyms</b>                         | TL1; VEG1   |
| <b>Host Species</b>                     | Rabbit  |
| <b>Description</b>                      | Biotinylated Anti-TNFSF15 antibody(DMC464); IgG1 Chimeric mAb   |
| <b>Delivery</b>                         | 2-3 weeks   |
| <b>Uniprot ID</b>                       | O95150  |
| <b>IgG type</b>                         | Rabbit/Human Fc chimeric IgG1   |
| <b>Clonality</b>                        | Monoclonal  |
| <b>Reactivity</b>                       | Human   |
| <b>Applications</b>                     | Flow Cyt  |
| <b>Recommended Dilutions</b>            | Flow Cyt 1:100  |
| <b>Purification</b>                     | Purified from cell culture supernatant by affinity chromatography   |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.  |
| <b>Storage&amp;Shipping</b>             | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| <b>Background</b>                       | The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This protein is abundantly expressed in endothelial cells; but is not expressed in either B or T cells. The expression of this protein is inducible by TNF and IL-1 alpha. This cytokine is a ligand for receptor TNFRSF25 and decoy receptor TNFRSF21:DR6. It can activate NF-kappaB and MAP kinases; and acts as an autocrine factor to induce apoptosis in endothelial cells. This cytokine is also found to inhibit endothelial cell proliferation; and thus may function as an angiogenesis inhibitor. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq; Feb 2011] |
| <b>Usage</b>                            | Research use only   |
| <b>Conjugate</b>                        | Biotinylated  |
| <b>DIMA Disclaimer</b>                  | All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.  |

