Cat. No. DMC100276B



PRODUCT INFORMATION

Clone ID **DMC276 Target VEGFA**

Synonyms MVCD1; VEGF; VPF

Host Species Rabbit

Biotinylated Anti-VEGFA antibody(DMC276); IgG1 Description

Chimeric mAb

Delivery 2-3 weeks **Uniprot ID** P15692

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended

Dilutions

Flow Cyt 1:100

Purified from cell culture supernatant by affinity **Purification**

chromatography

Formulation &

Background

Storage & Shipping

- 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Lyophilized from sterile PBS, pH 7.4. Normally 5 %

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene is a member of the PDGF:VEGF growth factor family. It encodes a heparin-binding protein; which exists as a disulfide-linked

homodimer. This growth factor induces proliferation and migration of vascular endothelial cells; and is essential for both physiological and pathological angiogenesis. Disruption of this gene in mice resulted in abnormal embryonic blood vessel formation. This gene is upregulated in many known tumors and its expression is correlated with tumor stage and progression. Elevated levels of this protein are found in patients with POEMS syndrome; also known as Crow-Fukase syndrome. Allelic variants of this gene have been associated with microvascular complications of diabetes 1 (MVCD1) and

atherosclerosis. Alternatively spliced transcript variants encoding different isoforms have been described. There is also evidence for alternative translation initiation from upstream non-AUG (CUG) codons resulting in additional isoforms. A recent study showed that a C-terminally extended isoform is produced by use of an alternative inframe translation termination codon via a stop codon readthrough mechanism; and that this isoform is antiangiogenic. Expression of some isoforms derived from the AUG start codon is regulated by a small upstream open reading frame; which is located within an internal

ribosome entry site.

Usage Research use only

Add Cersiugaten institute of Biotinylated Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com



Biotinylated Anti-VEGFA antibody(DMC276); IgG1 Chimeric mAb

Cat. No. DMC100276B



DIMA Disclaimer

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to JIMARBIO CONTRIBUTION CONTRIBUT ensure no IP infringement.

> Email: info@dimabio.com Website: www.dimabio.com



