

**PRODUCT INFORMATION**

<b>Target</b>	CDH17
<b>Synonyms</b>	HPT1; HPT-1
<b>Description</b>	Biotinylated Recombinant human CDH17 Protein with C-terminal His Avi tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q12864
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-6×His Avi Tag
<b>Molecular Characterization</b>	CDH17(Gln23-Met787) 6xHis Avi tag
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 87.5 kDa after removal of the signal peptide.
<b>Purity</b>	The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene is a member of the cadherin superfamily, genes encoding calcium-dependent, membrane-associated glycoproteins. The encoded protein is cadherin-like, consisting of an extracellular region, containing 7 cadherin domains, and a transmembrane region but lacking the conserved cytoplasmic domain. The protein is a component of the gastrointestinal tract and pancreatic ducts, acting as an intestinal proton-dependent peptide transporter in the first step in oral absorption of many medically important peptide-based drugs. The protein may also play a role in the morphological organization of liver and intestine. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2009]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Biotinylated






Figure 1. Biotinylated Human CDH17 Protein, His Tag on SDS-PAGE under reducing condition.

### Biotinylated Human CDH17, His Tagged protein ELISA

0.2  $\mu$ g of Anti-CDH17(MRG007 biosimilar) mAb per well




Figure 2. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Anti-CDH17(MRG007 biosimilar) mAb (BME100289) can bind Biotinylated Human CDH17 Protein, His Tag (PME100801B) in a linear range of 200-1000ng/mL.

