


## PRODUCT INFORMATION

<b>Tag</b>	C-Flag and Avi Tag
<b>Target</b>	CXCR7
<b>Synonyms</b>	RDC1; ACKR3; RDC-1; CMKOR1; CXCR7; GPR159
<b>Description</b>	Biotinylated Human CXCR7 full length protein-synthetic nanodisc
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P25106
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Transmembrane
<b>Protein Pathways</b>	Axon guidance, Chemokine signaling pathway, Cytokine-cytokine receptor interaction, Endocytosis, Leukocyte transendothelial migration
<b>Molecular Weight</b>	The human full length CXCR7 Protein has a MW of 44.4 kDa.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a member of the G-protein coupled receptor family. Although this protein was earlier thought to be a receptor for vasoactive intestinal peptide (VIP), it is now considered to be an orphan receptor, in that its endogenous ligand has not been identified. The protein is also a coreceptor for human immunodeficiency viruses (HIV). Translocations involving this gene and HMGA2 on chromosome 12 have been observed in lipomas. [provided by RefSeq, Jul 2008]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Biotinylated



**Biotinylated Human CXCR7 full length protein-synthetic nanodisc ELISA**

0.2 µg of Anti-Flag Rabbit mAb, Flag tagged protein per well




**Biotinylated Human CXCR7 full length protein-synthetic nanodisc (ng/mL)**

Figure 1. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Anti-Flag Rabbit mAb can bind Biotinylated Human CXCR7 full length protein-synthetic nanodisc(FLP100095B) in a linear range of 16-400 ng/mL. In order to specifically detect FLP100095B, HRP Conjugated Streptavidin was used as detection antibody.

**Biotinylated Human CXCR7 full length protein-synthetic nanodisc ELISA**

0.2 µg of Anti-CXCR7 antibody, IgG1 Chimeric mAb protein per well



**Biotinylated Human CXCR7 full length protein-synthetic nanodisc (ng/mL)**

Figure 2. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Anti-CXCR7 antibody, IgG1 Chimeric mAb can bind Biotinylated Human CXCR7 full length protein-synthetic nanodisc(FLP100095B) in a linear range of 16-400 ng/mL. In order to specifically detect FLP100095B, HRP Conjugated Streptavidin was used as detection antibody.



**Biotinylated Human CXCR7 full length protein-synthetic nanodisc ELISA**  
**0.2µg Biotinylated Human CXCR7-Nanodisc, Flag Tag per well**




Figure 3. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Biotinylated Human CXCR7 full length protein-synthetic nanodisc(FLP100095B) can bind Anti-Flag Rabbit mAb in a linear range of 0.64-16 ng/mL.





Figure 4. Biotinylated Human CXCR7-Nanodisc, Flag Tag on SDS-PAGE



**Biotinylated Human CXCR7 full length protein-synthetic nanodisc ELISA**

0.1 µg of Streptavidin per well



**Biotinylated Human CXCR7 full length protein-synthetic nanodisc.(ng/ml)**

Figure 5. ELISA plate pre-coated by 1 µg/mL (100 µL/well) Streptavidin can bind Biotinylated Human CXCR7 full length protein-synthetic nanodisc(FLP100095B) in a linear range of 16-400 ng/mL. In order to specifically detect FLP100095B, Anti-Flag Rabbit antibody was used as detection antibody.

