

PRODUCT INFORMATION

Tag	C-Flag&Avi Tag
Expression Host	HEK293
Target	NK2R
Synonyms	TACR2, NKNAR, SKR, TAC2R
Description	Biotinylated Human NK2R full length protein-synthetic nanodisc
Uniprot ID	P21452
Protein Families	Transmembrane
Protein Pathways	Peptide GPCRs,
Molecular Weight	The human full length NK2R protein has a MW of 49.2kDa
Delivery	In Stock
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene belongs to a family of genes that function as receptors for tachykinins. Receptor affinities are specified by variations in the 5'-end of the sequence. The receptors belonging to this family are characterized by interactions with G proteins and 7 hydrophobic transmembrane regions. This gene encodes the receptor for the tachykinin neuropeptide substance K, also referred to as neurokinin A. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Biotinylated







Figure 1. Biotinylated Human NK2R-Nanodisc, Flag Tag on SDS-PAGE

Biotinylated Human NK2R full length protein-synthetic nanodisc ELISA

0.1 μg of Streptavidin protein per well



Biotinylated Human NK2R full length protein-synthetic nanodisc (ng/mL)

Figure 2. On a streptavidin-coated plate (0.1 μg /well) immobilized with biotinylated Human NK2R full-length protein-synthetic nanodisc (FLP100360B, 1 $\mu\text{g}/\text{mL}$, 100 $\mu\text{L}/\text{well}$), the Anti-Flag Rabbit mAb showed linear binding in the range of 0.64-16 ng/mL. In order to specifically detect FLP100360B, HRP Conjugated Goat anti Rabbit was used as detection antibody.






Figure 3. WB analysis of Human NK2R-Nanodisc with Streptavidin HRP at 1/4000 dilution

DIMABIO CONFIDENTIAL

