

PRODUCT INFORMATION

Target	NKG2D
Synonyms	KLR; CD314; KLRK1; NKG2-D; D12S2489E
Description	Biotinylated Recombinant human NKG2D Protein with N-terminal Avi Human Fc tag
Delivery	In Stock
Uniprot ID	P26718
Expression Host	HEK293
Tag	N-Avi Human Fc tag
Molecular Characterization	Avi hFc(Glu99-Ala330)tag NKG2D(Ile73-Val216)
Molecular Weight	The protein has a predicted molecular mass of 44.6 kDa after removal of the signal peptide.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Natural killer (NK) cells are lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. NK cells preferentially express several calcium-dependent (C-type) lectins, which have been implicated in the regulation of NK cell function. The NKG2 gene family is located within the NK complex, a region that contains several C-type lectin genes preferentially expressed in NK cells. This gene encodes a member of the NKG2 family. The encoded transmembrane protein is characterized by a type II membrane orientation (has an extracellular C terminus) and the presence of a C-type lectin domain. It binds to a diverse family of ligands that include MHC class I chain-related A and B proteins and UL-16 binding proteins, where ligand-receptor interactions can result in the activation of NK and T cells. The surface expression of these ligands is important for the recognition of stressed cells by the immune system, and thus this protein and its ligands are therapeutic targets for the treatment of immune diseases and cancers. Read-through transcription exists between this gene and the upstream KLRC4 (killer cell lectin-like receptor subfamily C, member 4) family member in the same cluster. [provided by RefSeq, Dec 2010]
Usage	Research use only
Conjugate	Biotinylated






Figure 1. Biotinylated Human NKG2D Protein, hFc Tag on SDS-PAGE under reducing condition.

Biotinylated Human NKG2D, hFc Tagged Protein ELISA

0.2 µg of Anti-NKG2D(5C5 biosimilar) mAb per well




Figure 2. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Anti-NKG2D(5C5 biosimilar) mAb (BME100207) can bind Biotinylated Human NKG2D Protein, hFc Tag (PME101956B) in a linear range of 200-1000 ng/mL.

