

**PRODUCT INFORMATION**

<b>Target</b>	ANXA2
<b>Synonyms</b>	Annexin
<b>Description</b>	Recombinant Cynomolgus ANXA2 protein with C-terminal 6×His tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	A0A2K5UQS9
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-6×His Tag
<b>Molecular Characterization</b>	ANXA2(Ser2-Asp339) 6×His tag
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 39.3 kDa after removal of the signal peptide. The apparent molecular mass of cANXA2-His is approximately 35-55 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a member of the annexin family. Members of this calcium-dependent phospholipid-binding protein family play a role in the regulation of cellular growth and in signal transduction pathways. This protein functions as an autocrine factor which heightens osteoclast formation and bone resorption. This gene has three pseudogenes located on chromosomes 4, 9 and 10, respectively. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this gene. Annexin A2 expression has been found to correlate with resistance to treatment against various cancer forms. [provided by RefSeq, Dec 2019]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated






Figure 1. Cynomolgus ANXA2 Protein, His Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

