

PRODUCT INFORMATION

CCR8 **Target**

Synonyms Chemokine receptor 8

Recombinant Cynomolgus CCR8 protein with C-Description

terminal human Fc tag

Delivery In Stock **Uniprot ID** G7NYJ2 **HEK293 Expression Host**

Tag C-Human Fc Tag

Molecular

Storage & Shipping

Background

CCR8(Met1-Lys35) hFc(Glu99-Ala330) Characterization

The protein has a predicted molecular mass of **Molecular Weight**

30.2 kDa after removal of the signal peptide. The apparent molecular mass of cCCR8-hFc is

approximately 35-55 kDa due to glycosylation. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

Purity staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & Reconstitution

 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a member of the beta chemokine receptor family, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. Chemokines and their receptors are important for the migration of various cell types into the inflammatory sites. This receptor protein preferentially expresses in the thymus. I-309, thymus activation-regulated

cytokine (TARC) and macrophage inflammatory protein-1 beta (MIP-1 beta) have been identified as ligands of this receptor. Studies of this receptor and its ligands suggested its role in regulation of monocyte chemotaxis and thymic cell apoptosis. More specifically, this receptor may contribute to the proper positioning of activated T cells within the antigenic challenge sites and specialized areas of lymphoid tissues. This gene is located at the chemokine receptor gene cluster region. [provided by RefSeq, Jul 2008]

> Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)






Figure 1. Cynomolgus CCR8 Protein, hFc Tag on SDS-PAGE under reducing condition.

