

PRODUCT INFORMATION

Target	CD47
Synonyms	IAP; OA3; MER6
Description	Recombinant Cynomolgus CD47 protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	A0A2K5X4H8
Expression Host	HEK293
Tag	C-Human Fc tag
Molecular Characterization	CD47(Gln19-Glu141) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 40.0 kDa after removal of the signal peptide. The apparent molecular mass of cCD47-hFc is approximately 55-100 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
Background	This gene encodes a membrane protein, which is involved in the increase in intracellular calcium concentration that occurs upon cell adhesion to extracellular matrix. The encoded protein is also a receptor for the C-terminal cell binding domain of thrombospondin, and it may play a role in membrane transport and signal transduction. This gene has broad tissue distribution, and is reduced in expression on Rh erythrocytes. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2010]
Usage	Research use only
Conjugate	Unconjugated





Figure 1. Cynomolgus CD47 Protein, hFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

