

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	5HT1B
Synonyms	5-HT-1B; 5-HT-1D-beta; 5-HT1B; 5-HT1DB; HTR1D2; HTR1DB; S12
Description	Human 5HT1B-Strep full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P28222
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Neuroactive ligand-receptor interaction
Molecular Weight	The human full length 5HT1B-Strep protein has a MW of 43.6 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	A G-protein coupled receptor for serotonin (5-hydroxytryptamine). Ligand binding activates second messengers that inhibit the activity of adenylate cyclase and manage the release of serotonin, dopamine, and acetylcholine in the brain. The encoded protein may be involved in several neuropsychiatric disorders and therefore is often a target of antidepressant and other psychotherapeutic drugs.
Usage	Research use only
Conjugate	Unconjugated




ELISA assay to evaluate 5HT1B-Strep-Nanodisc
0.2 μ g Human 5HT1B-Strep-Nanodisc per well

Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag 5HT1B-Strep-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with 5HT1B-Strep-nanodisc is 7.699ng/ml.




Figure 2. Human 5HT1B-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

