

## PRODUCT INFORMATION

<b>Tag</b>	C-Flag Tag
<b>Target</b>	ABCG1
<b>Synonyms</b>	ABC8; WHITE1
<b>Description</b>	Human ABCG1 full length protein-synthetic nanodisc
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P45844
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Druggable Genome, Transmembrane
<b>Protein Pathways</b>	ABC transporters
<b>Molecular Weight</b>	The human full length ABCG1 protein has a MW of 75.6 kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
<b>Background</b>	The protein is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the White subfamily. It is involved in macrophage cholesterol and phospholipids transport, and may regulate cellular lipid homeostasis in other cell types
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated



### ELISA assay to evaluate ABCG1-Nanodisc 0.2 $\mu$ g Human ABCG1-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag ABCG1-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with ABCG1-Nanodisc is 9.545ng/ml.



Figure2. Human ABCG1-Nanodisc, Flag Tag on SDS-PAGE

