

## **PRODUCT INFORMATION**

ACE2 **Target** 

**Synonyms** ACE-2; ACEH; ACE2

Recombinant Human ACE2 Protein with N-**Description** 

terminal 6×His tag

**Delivery** In Stock **Uniprot ID** Q9BYF1 **Expression Host HEK293** Tag N-6×His Tag

Molecular

Storage & Shipping

**Background** 

6×His ACE2(Gln18-Ser740) Characterization

The protein has a predicted molecular mass of

84.4 kDa after removal of the signal peptide. The apparent molecular mass of His-ACE2 is **Molecular Weight** 

approximately 100-130 kDa due to glycosylation.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene belongs to the angiotensin-converting enzyme family of dipeptidyl carboxydipeptidases and has

considerable homology to human angiotensin 1 converting enzyme. This secreted protein catalyzes the cleavage of angiotensin I into angiotensin 1-9, and angiotensin II into the vasodilator angiotensin 1-7. The organ- and cell-

specific expression of this gene suggests that it may play a role in the regulation of cardiovascular and renal function, as well as fertility. In addition, the encoded protein is a functional receptor for the spike glycoprotein of the human coronaviruses SARS and HCoV-NL63.

> Email: info@dimabio.com Website: www.dimabio.com

**Usage** Research use only Conjugate Unconjugated








Figure 1. Human ACE2 Protein, His Tag on SDS-PAGE under reducing condition.

## Human ACE2, His Tagged protein ELISA

0.2 µg of ACE2, His Tagged protein per well




Figure 2. ELISA plate pre-coated by 2  $\mu$ g/ml (100  $\mu$ l/well) Human ACE2, His Tagged protein PME100490 can bind S-RBD, hFc tagged protein PME100487 in a linear range of 0.128-80.0 ng/ml.

