

## PRODUCT INFORMATION

<b>Target</b>	ACE2
<b>Synonyms</b>	ACE-2;ACEH;ACE2
<b>Description</b>	Recombinant Human ACE2 Protein with N-terminal 6×His tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q9BYF1
<b>Expression Host</b>	HEK293
<b>Tag</b>	N-6×His Tag
<b>Molecular Characterization</b>	6×His ACE2(Gln18-Ser740)
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 84.4 kDa after removal of the signal peptide. The apparent molecular mass of His-ACE2 is approximately 100-130 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein encoded by this gene belongs to the angiotensin-converting enzyme family of dipeptidyl carboxydipeptidases and has considerable homology to human angiotensin 1 converting enzyme. This secreted protein catalyzes the cleavage of angiotensin I into angiotensin 1-9, and angiotensin II into the vasodilator angiotensin 1-7. The organ- and cell-specific expression of this gene suggests that it may play a role in the regulation of cardiovascular and renal function, as well as fertility. In addition, the encoded protein is a functional receptor for the spike glycoprotein of the human coronaviruses SARS and HCoV-NL63.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated






Figure 1. Human ACE2 Protein, His Tag on SDS-PAGE under reducing condition.

### Human ACE2, His Tagged protein ELISA

0.2 µg of ACE2, His Tagged protein per well




Figure 2. ELISA plate pre-coated by 2 µg/ml (100 µl/well) Human ACE2, His Tagged protein [PME100490] can bind S-RBD, hFc tagged protein PME100487 in a linear range of 0.128-80.0 ng/ml.

