

**PRODUCT INFORMATION**

<b>Tag</b>	C-Flag&Strep Tag
<b>Expression Host</b>	HEK293
<b>Target</b>	ACHB
<b>Synonyms</b>	ACHRB, CHRNB, CMS1D, CMS2A, CMS2C, SCCMS
<b>Description</b>	Human ACHB-Strep full length protein-synthetic nanodisc
<b>Uniprot ID</b>	P11230
<b>Protein Families</b>	Ion Channels: Cys-loop Receptors
<b>Protein Pathways</b>	N/A
<b>Molecular Weight</b>	The human full length ACHB-Strep protein has a MW of 56.7 kDa
<b>Delivery</b>	6~8weeks
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The muscle acetylcholine receptor is composed of five subunits: two alpha subunits and one beta, one gamma, and one delta subunit. This gene encodes the beta subunit of the acetylcholine receptor. The acetylcholine receptor changes conformation upon acetylcholine binding leading to the opening of an ion-conducting channel across the plasma membrane. Mutations in this gene are associated with slow-channel congenital myasthenic syndrome. [provided by RefSeq, Jul 2008]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated

