

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Expression Host	HEK293
Target	ACM1
Synonyms	HM1, M1, M1R
Description	Human ACM1-Strep full length protein-synthetic nanodisc
Uniprot ID	P11229
Protein Families	GPCR,Transmembrane,Druggable Genome,
Protein Pathways	Calcium regulation in cardiac cells,GPCRDB Class A Rhodopsin-like,Monoamine GPCRs,Regulation of Actin Cytoskeleton KEGG,
Molecular Weight	The human full length ACM1-Strep protein has a MW of 51.4 kDa
Delivery	6~8weeks
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The muscarinic cholinergic receptors belong to a larger family of G protein-coupled receptors. The functional diversity of these receptors is defined by the binding of acetylcholine and includes cellular responses such as adenylate cyclase inhibition, phosphoinositide degeneration, and potassium channel mediation. Muscarinic receptors influence many effects of acetylcholine in the central and peripheral nervous system. The muscarinic cholinergic receptor 1 is involved in mediation of vagally-induced bronchoconstriction and in the acid secretion of the gastrointestinal tract. The gene encoding this receptor is localized to 11q13. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated

