

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	ACM3
Synonyms	EGBRS, HM3, PBS
Description	Human ACM3 full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	P20309
Expression Host	HEK293
Protein Families	GPCR,Transmembrane,Druggable Genome, Calcium regulation in cardiac cells, GPCRDB Class A Rhodopsin-like, GPCRDB Other, Monoamine GPCRs, Regulation of Actin Cytoskeleton KEGG, Metabolic and Obesity
Protein Pathways	
Molecular Weight	The human full length ACM3 protein has a MW of 66.1kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The muscarinic cholinergic receptors belong to a larger family of G protein-coupled receptors. The functional diversity of these receptors is defined by the binding of acetylcholine and includes cellular responses such as adenylate cyclase inhibition, phosphoinositide degeneration, and potassium channel mediation. Muscarinic receptors influence many effects of acetylcholine in the central and peripheral nervous system. The muscarinic cholinergic receptor 3 controls smooth muscle contraction and its stimulation causes secretion of glandular tissue. Alternative promoter use and alternative splicing results in multiple transcript variants that have different tissue specificities. [provided by RefSeq, Dec 2016]
Usage	Research use only
Conjugate	Unconjugated

