

PRODUCT INFORMATION

Target ADRB1

Synonyms ADRB1R; B1AR; BETA1AR; FNSS2; RHR

Human ADRB1 full length protein-synthetic **Description**

nanodisc

In Stock Delivery **Uniprot ID** P08588 **Expression Host HEK293**

Storage & Shipping

Background

Protein Families Druggable Genome, GPCR, Transmembrane

Calcium signaling pathway, Dilated cardiomyopathy, Endocytosis, Gap junction, **Protein Pathways**

Neuroactive ligand-receptor interaction

The human full length ADRB1 protein has a MW of **Molecular Weight**

51.2 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% Formulation & 8% trehalose is added as protectants before Reconstitution lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments. Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The adrenergic receptors (subtypes alpha 1, alpha 2, beta 1, and beta 2) are a prototypic family of guanine nucleotide binding regulatory protein-coupled receptors that mediate the

physiological effects of the hormone epinephrine and the neurotransmitter norepinephrine. Beta-1 adrenoceptors are predominately located in the heart. Specific polymorphisms in this gene have been shown to affect the resting heart rate and

> Email: info@dimabio.com Website: www.dimabio.com

can be involved in heart failure.

Usage Research use only

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)





ELISA assay to evaluate ADRB1-Nanodisc 0.2µg Human ADRB1-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag ADRB1-Nanodisc ($0.2\mu g/per$ well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with ADRB1-Nanodisc is 6.812ng/ml.



Figure 2. Human ADRB1-Nanodisc, Flag Tag on SDS-PAGE

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

