

## PRODUCT INFORMATION

|   |   |
|---|---|
| <b>Tag</b>                              | C-Flag Tag  |
| <b>Target</b>                           | ADRB2   |
| <b>Synonyms</b>                         | ADRB2R, ADRBR, B2AR, BAR, BETA2AR   |
| <b>Description</b>                      | Human ADRB2 full length protein-synthetic nanodisc  |
| <b>Delivery</b>                         | 6~8weeks  |
| <b>Uniprot ID</b>                       | P07550  |
| <b>Expression Host</b>                  | HEK293  |
| <b>Protein Families</b>                 | GPCR,Transmembrane,Druggable Genome,<br>Calcium regulation in cardiac cells,GPCRDB Class A Rhodopsin-like,GPCRDB Other,Monoamine  |
| <b>Protein Pathways</b>                 | GPCRs,Metabolic and Obesity,G-Protein Coupled Receptors Signaling Pathway,cAMP and Ca <sup>2+</sup> Signaling Pathway,  |
| <b>Molecular Weight</b>                 | The human full length ADRB2 protein has a MW of 46.5kDa   |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.  |
| <b>Storage&amp;Shipping</b>             | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| <b>Background</b>                       | This gene encodes beta-2-adrenergic receptor which is a member of the G protein-coupled receptor superfamily. This receptor is directly associated with one of its ultimate effectors, the class C L-type calcium channel Ca(V)1.2. This receptor-channel complex also contains a G protein, an adenylyl cyclase, cAMP-dependent kinase, and the counterbalancing phosphatase, PP2A. The assembly of the signaling complex provides a mechanism that ensures specific and rapid signaling by this G protein-coupled receptor. This receptor is also a transcription regulator of the alpha-synuclein gene, and together, both genes are believed to be associated with risk of Parkinson's Disease. This gene is intronless. Different polymorphic forms, point mutations, and/or downregulation of this gene are associated with nocturnal asthma, obesity, type 2 diabetes and cardiovascular disease. [provided by RefSeq, Oct 2019] |
| <b>Usage</b>                            | Research use only   |
| <b>Conjugate</b>                        | Unconjugated  |

