

PRODUCT INFORMATION

ALK **Target**

Synonyms CD246;NBLST3

Recombinant Human ALK Protein with C-terminal **Description**

human Fc tag

Delivery In Stock **Uniprot ID** Q9UM73 **Expression Host** HFK293

Tag C-Human Fc Tag

Molecular

Purity

Background

ALK(Val19-Ser1038) hFc(Glu99-Ala330) Characterization

The protein has a predicted molecular mass of

136.4 kDa after removal of the signal peptide. The apparent molecular mass of ALK-hFc is **Molecular Weight**

approximately 130-250 kDa due to glycosylation.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation &

- 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a receptor tyrosine kinase, which belongs to the insulin receptor superfamily. This protein comprises an extracellular domain, an hydrophobic stretch corresponding to a single pass transmembrane region, and an intracellular kinase domain. It plays an important role in the development of the brain and exerts its effects on specific neurons in the nervous system. This gene

has been found to be rearranged, mutated, or amplified in a series of tumours including anaplastic large cell lymphomas, neuroblastoma, and non-small cell lung cancer. The chromosomal

rearrangements are the most common genetic alterations in this gene, which result in creation of multiple fusion genes in tumourigenesis, including ALK (chromosome 2)/EML4 (chromosome 2), ALK/RANBP2 (chromosome 2), ALK/ATIC

(chromosome 2), ALK/TFG (chromosome 3), ALK/NPM1 (chromosome 5), ALK/SQSTM1 (chromosome 5), ALK/KIF5B (chromosome 10), ALK/CLTC (chromosome 17), ALK/TPM4 (chromosome 19), and ALK/MSN1 (chromosome

> Email: info@dimabio.com Website: www.dimabio.com

X).[provided by RefSeq, Jan 2011]

Usage Research use only

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)






Figure 1. Human ALK Protein, hFc Tag on SDS-PAGE under reducing condition



