

PRODUCT INFORMATION

ANPRB **Target**

AMDM;ANPb;NPR2;ECDM;GC-B;GCB;GUC2B;GUCY2B;NPRB;NPRBi;SNSK **Synonyms**

Description Recombinant Human ANPRB Protein with C-terminal 6×His tag

Delivery P20594 **Uniprot ID Expression Host HEK293** C-6×His Tag

Molecular Characterization ANPRB(Arg23-Ile458) 6×His tag

The protein has a predicted molecular mass of 49.3 kDa after removal **Molecular Weight**

of the signal peptide.

The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining. **Purity**

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Store at -20°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized Storage & Shipping

proteins are shipped at ambient temperature.

proteins are shipped at ambient temperature.

This gene encodes natriuretic peptide receptor B, one of two integral membrane receptors for natriuretic peptides. Both NPR1 and NPR2 contain five functional domains: an extracellular ligand-binding domain, a single membrane-spanning region, and intracellularly a protein kinase homology domain, a helical hinge region involved in oligomerization, and a carboxyl-terminal guanylyl cyclase catalytic domain. The protein is the primary receptor for C-type natriuretic peptide (CNP), which upon ligand binding exhibits greatly increased guanylyl cyclase activity. Mutations in this gene are the cause of acromesomelic dysplasia Maroteaux type. [provided by RefSeq, Jul 2008]

Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only Conjugate Unconjugated

Background








Figure 1. Human ANPRB Protein, His Tag on SDS-PAGE under reducing condition.



