

PRODUCT INFORMATION

Tag	C-Flag Tag
Expression Host	HEK293
Target	AQP5
Synonyms	AQP-5; PPKB
Description	Human AQP5 full length protein-synthetic nanodisc
Uniprot ID	P55064
Protein Families	Druggable Genome, Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length AQP5 protein has a MW of 28.3 kDa
Delivery	In Stock
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Aquaporin 5 (AQP5) is a water channel protein. Aquaporins are a family of small integral membrane proteins related to the major intrinsic protein (MIP or AQP0). Aquaporin 5 plays a role in the generation of saliva, tears and pulmonary secretions. AQP0, AQP2, AQP5, and AQP6 are closely related and all map to 12q13.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate AQP5-Nanodisc 0.2 μ g Human AQP5-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag AQP5-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with AQP5-Nanodisc is 5.102ng/ml.




Figure2. Human AQP5-Nanodisc, Flag Tag on SDS-PAGE

