

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	ASIC3
Synonyms	ACCN3, DRASIC, SLNAC1, TNaC1
Description	Human ASIC3 full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q9UHC3
Expression Host	HEK293
Protein Families	Ion Channels: Other
Protein Pathways	N/A
Molecular Weight	The human full length ASIC3 protein has a MW of 58.9kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a member of the degenerin/epithelial sodium channel (DEG/ENaC) superfamily. The members of this family are amiloride-sensitive sodium channels that contain intracellular N and C termini, two hydrophobic transmembrane regions, and a large extracellular loop, which has many cysteine residues with conserved spacing. The member encoded by this gene is an acid sensor and may play an important role in the detection of lasting pH changes. In addition, a heteromeric association between this member and acid-sensing (proton-gated) ion channel 2 has been observed as proton-gated channels sensitive to gadolinium. Alternatively spliced transcript variants have been described. [provided by RefSeq, Feb 2012]
Usage	Research use only
Conjugate	Unconjugated

