Cat. No. FLP100031A



## **PRODUCT INFORMATION**

Tag C-Flag Tag

Target GPR75

**Synonyms** GPRchr2; WI31133

**Description**Human beta2AR-GPR75-BRIL full length protein-

synthetic nanodisc

Delivery In Stock
Uniprot ID 095800
Expression Host HEK293

**Protein Families** Druggable Genome, GPCR, Transmembrane

Protein Pathways N/A

Molecular Weight
The human full length beta2AR-GPR75-BRIL

protein has a MW of 59.4 kDa

Formulation & Reconstitution

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

**Storage & Shipping** intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

GPR75 is a member of the G protein-coupled receptor family and is a novel target for the clinical treatment of obesity. Human GPR75 haploinsufficiency exhibits a striking phenotype of low body fat, and GPR75 knockout mice are hypophagic and thin, improving glucose tolerance and insulin sensitivity. To enhance the expression level of GPR75, twenty-four amino acids from the

Background beta2AR receptor

(MGQPGNGSAFLLAPNRSHAPDHDV) were used to replace the original residues1-31 in a modified version of the GPR75 receptor. Furthermore, the BRIL sequence was used to replace the original ICL3 loop (237-306) to enhance receptor stability. BRIL is thermostabilized apocytochrome b562 with mutations M7W, H102I, R106L, PDB ID 1M6T.

Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only
Conjugate Unconjugated



Cat. No. FLP100031A



## ELISA assay to evaluate β2AR-GPR75-BRIL-Nanodisc 0.2μg Human β2AR-GPR75-BRIL-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag beta 2AR-GPR75-BRIL-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with beta 2AR-GPR75-BRIL-Nanodisc is 4.678 ng/ml.




Figure 2. Human beta2AR-GPR75-BRIL-Nanodisc, Flag Tag on SDS-PAGE



Email: info@dimabio.com Website: www.dimabio.com

