

PRODUCT INFORMATION

| | |
|---|--|
| Target | BAFF |
| Synonyms | DTL; TNFSF13B; BLYS; CD257; TALL1; THANK; ZTNF4; TALL-1; TNLG7A; TNFSF20 |
| Description | Recombinant human BAFF Protein with N-terminal mouse Fc tag and C-terminal 6xHis tag |
| Delivery | In Stock |
| Uniprot ID | Q9Y275 |
| Expression Host | HEK293 |
| Tag | N-Mouse Fc tag and C-6xHis tag |
| Molecular Characterization | mFc(Pro99-Lys330) BAFF(Leu72-Leu285) 6xHis tag |
| Molecular Weight | The protein has a predicted molecular mass of 51.8 kDa after removal of the signal peptide. |
| Purity | The purity of the protein is greater than 90% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage&Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for receptors TNFRSF13B/TACI, TNFRSF17/BCMA, and TNFRSF13C/BAFFR. This cytokine is expressed in B cell lineage cells, and acts as a potent B cell activator. It has been also shown to play an important role in the proliferation and differentiation of B cells. Alternatively spliced transcript variants encoding distinct isoforms have been identified. [provided by RefSeq, Mar 2011] |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. Human BAFF Protein, mFc-His Tag on SDS-PAGE under reducing condition.

