

## PRODUCT INFORMATION

<b>Target</b>	BAFF
<b>Synonyms</b>	DTL; TNFSF13B; BLYS; CD257; TALL1; THANK; ZTNF4; TALL-1; TNLG7A; TNFSF20
<b>Description</b>	Recombinant human BAFF Protein with N-terminal mouse Fc tag and C-terminal 6×His tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q9Y275
<b>Expression Host</b>	HEK293
<b>Tag</b>	N-Mouse Fc tag and C-6×His tag
<b>Molecular Characterization</b>	mFc(Pro99-Lys330) BAFF(Leu72-Leu285) 6×His tag
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 51.8 kDa after removal of the signal peptide.
<b>Purity</b>	The purity of the protein is greater than 90% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for receptors TNFRSF13B/TACI, TNFRSF17/BCMA, and TNFRSF13C/BAFFR. This cytokine is expressed in B cell lineage cells, and acts as a potent B cell activator. It has been also shown to play an important role in the proliferation and differentiation of B cells. Alternatively spliced transcript variants encoding distinct isoforms have been identified. [provided by RefSeq, Mar 2011]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated






Figure 1. Human BAFF Protein, mFc-His Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

