

PRODUCT INFORMATION

|                              |  |
|------------------------------|--|
| Target                       | BAI1   |
| Synonyms                     | ADGRB1; GDAIF  |
| Description                  | Recombinant human BAI1(31-926) Protein with C-terminal Flag and Strep Tag  |
| Delivery                     | In Stock   |
| Uniprot ID                   | O14514   |
| Expression Host              | HEK293   |
| Tag                          | C-Flag and Strep Tag   |
| Molecular Characterization   | BAI1(Ala31-Leu926) Flag and Strep Tag  |
| Molecular Weight             | The protein has a predicted molecular mass of 102.5 kDa after removal of the signal peptide.   |
| Purity                       | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.   |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.   |
| Storage&Shipping             | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.  |
| Background                   | Angiogenesis is controlled by a local balance between stimulators and inhibitors of new vessel growth and is suppressed under normal physiologic conditions. Angiogenesis has been shown to be essential for growth and metastasis of solid tumors. In order to obtain blood supply for their growth, tumor cells are potently angiogenic and attract new vessels as results of increased secretion of inducers and decreased production of endogenous negative regulators. BAI1 contains at least one 'functional' p53-binding site within an intron, and its expression has been shown to be induced by wildtype p53. There are two other brain-specific angiogenesis inhibitor genes, designated BAI2 and BAI3 which along with BAI1 have similar tissue specificities and structures, however only BAI1 is transcriptionally regulated by p53. BAI1 is postulated to be a member of the secretin receptor family, an inhibitor of angiogenesis and a growth suppressor of glioblastomas [provided by RefSeq, Jul 2008] |
| Usage                        | Research use only  |
| Conjugate                    | Unconjugated   |






Figure 1. Human BAI1(31-926) Protein, Flag and Strep Tag on SDS-PAGE under reducing condition.

