

**PRODUCT INFORMATION**

|   |  |
|---|--|
| <b>Tag</b>                              | C-Flag Tag   |
| <b>Expression Host</b>                  | HEK293   |
| <b>Target</b>                           | CA2D4  |
| <b>Synonyms</b>                         | RCD4   |
| <b>Description</b>                      | Human CA2D4 full length protein-synthetic nanodisc   |
| <b>Uniprot ID</b>                       | Q7Z3S7   |
| <b>Protein Families</b>                 | Ion Channels: Other  |
| <b>Protein Pathways</b>                 | N/A  |
| <b>Molecular Weight</b>                 | The human full length CA2D4 protein has a MW of 127.9kDa   |
| <b>Delivery</b>                         | 6~8weeks   |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution  |
| <b>Sterility</b>                        | Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.  |
| <b>Storage&amp;Shipping</b>             | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.  |
| <b>Background</b>                       | This gene encodes a member of the alpha-2/delta subunit family, a protein in the voltage-dependent calcium channel complex. Calcium channels mediate the influx of calcium ions into the cell upon membrane polarization and consist of a complex of alpha-1, alpha-2/delta, beta, and gamma subunits in a 1:1:1:1 ratio. Various versions of each of these subunits exist, either expressed from similar genes or the result of alternative splicing. Research on a highly similar protein in rabbit suggests the protein described in this record is cleaved into alpha-2 and delta subunits. Alternate transcriptional splice variants of this gene have been observed but have not been thoroughly characterized. [provided by RefSeq, Jul 2008] |
| <b>Usage</b>                            | Research use only  |
| <b>Conjugate</b>                        | Unconjugated   |

