

PRODUCT INFORMATION

Tag	C-Flag Tag
Expression Host	HEK293
Target	CAC1A
Synonyms	APCA, BI, CACNL1A4, CAV2.1, DEE42, EA2, EIEE42, FHM, HPCA, MHP, MHP1, SCA6
Description	Human CAC1A full length protein-synthetic nanodisc
Uniprot ID	O00555
Protein Families	Ion Channels: Calcium
Protein Pathways	N/A
Molecular Weight	The human full length CAC1A protein has a MW of 282.6kDa
Delivery	6~8weeks
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Voltage-dependent calcium channels mediate the entry of calcium ions into excitable cells, and are also involved in a variety of calcium-dependent processes, including muscle contraction, hormone or neurotransmitter release, and gene expression. Calcium channels are multisubunit complexes composed of alpha-1, beta, alpha-2/delta, and gamma subunits. The channel activity is directed by the pore-forming alpha-1 subunit, whereas, the others act as auxiliary subunits regulating this activity. The distinctive properties of the calcium channel types are related primarily to the expression of a variety of alpha-1 isoforms, alpha-1A, B, C, D, E, and S. This gene encodes the alpha-1A subunit, which is predominantly expressed in neuronal tissue. Mutations in this gene are associated with 2 neurologic disorders, familial hemiplegic migraine and episodic ataxia 2. This gene also exhibits polymorphic variation due to (CAG) _n -repeats. Multiple transcript variants encoding different isoforms have been found for this gene. In one set of transcript variants, the (CAG) _n -repeats occur in the 3' UTR, and are not associated with any disease. But in another set of variants, an insertion extends the coding region to include the (CAG) _n -repeats which encode a polyglutamine tract. Expansion of the (CAG) _n -repeats from the normal 4-18 to 21-33 in the coding region is associated with spinocerebellar ataxia 6. [provided by RefSeq, Jul 2016]
Usage	Research use only
Conjugate	Unconjugated



DIMABIO CONFIDENTIAL

